

Monitoring Volcanoes

using the
***iGrav*TM Superconducting Gravity Meter**



*iGrav*TM Provides:

- ✓ Continuous Gravity Data
- ✓ Sub- μ Gal Precision
- ✓ Ultra-low Linear Drift
- ✓ Extremely Low Noise
- ✓ Constant Scale Factor
- ✓ Insensitive to Local Environment
- ✓ Tilt Correction

High sensitivity, allows measuring typical volcanic processes as far as 30 km away from the edifice:

- ✓ Safe distance
- ✓ Discriminating between shallow and deeper processes

Introduction

Simultaneous Gravity and GPS measurements have been proven a powerful combination for detecting subsurface mass or density changes long before eruption precursors appear. High precision continuous gravity measurements are paramount for detecting fast changes in density and phase, as well as magma movements and separating seasonal effects from deeper processes that may occur over years. Nonetheless, the standard spring gravity meters used for these measurements are limited to 10 to 15 μGal precision and are degraded by non-linear drifts, scale factor variability and environmental effects. These problems interfere with modeling volcanic structural changes taking place over years to decades.

GWR Instruments, Inc. is introducing the *iGrav* Superconducting Gravity Meter as an ideal choice for continuous volcano monitoring. The *iGrav* will improve precision to sub- μGal levels and will eliminate drifts and scale factor variability. The *iGrav* will provide a high precision, continuous gravity record that is ideal for interpreting volcanic activity and correlating with other geophysical measurements.

Detection of Mass Redistribution Long Before Eruption Precursors Appear

➤ Magma Movements on Different Time Scales:

- Intrusive mechanism leading to eruption
- Magma drainage out of a reservoir, magma migration from central conduit to fractures
- Magma storage
- Accumulation of magma at depth, increasing pressurization in the volcanic conduit and potential for explosive eruption
- Gas release
- Vesiculation (density decrease) and crystallization (density increase)
- Monitoring mass changes without significant vertical deformation (e.g. increase in CO_2 increases magma compressibility (Bonvalot et al., 2008))
- The *iGrav* is an indispensable forecasting tool for monitoring aseismic mass changes (e.g. Campi Flegrei or Phlegraean Fields, Naples)

➤ Volcano-groundwater Interactions:

- Discriminating between the intrusion of magma, water and gas
- Groundwater dynamics within a volcanic edifice (potential for slope failure, lahar generation and phreatic eruptions)
- Hydrothermal systems (e.g. Long Valley and Yellowstone calderas, Mt Uzu, Mt Sakurajima)

➤ Continuous Gravity Monitoring Enables:

- Avoiding aliasing from intermittent 4D gravity campaigns
- Recovering functioning laws and therefore, discriminating clearly between models for sources of volcano unrest
- Investigating short period (e.g. bubble formation and collapse in Strombolian activity) gravity changes
- Development of an accurate surface hydrological model to achieve μGal precision
- Precise modeling of Earth and ocean tides and atmospheric signals
- Changes in the mechanical response of the edifice to the tidal forces
- Providing a precise reference point to anchor microgravity surveys

➤ Integration and Correlation with Other Techniques:

- Combining with GPS – Most powerful combination to separate and interpret change in gravity with expansion, deflation or stability of volcanoes' surface
- Combining with Absolute Gravity extends spectrum of results to years and decades for studying long term reservoir activity
- Correlation with seismic activity and long period seismic signals

Network of *iGravs*


- Allow Monitoring at Safe Distance
- Enable Discriminating Between Shallow and Deep Processes
- Horizontal Gradients Probe Mass / Density Variations at Depth Over Both Space and Time
 - Two *iGravs* at distances of 1 and 10 km may allow discriminating between point sources from dyke intrusion or between shallow and deeper processes
 - One *iGrav* can measure the influence of a 2 x 3000 m dyke or of a 10^{12} kg point source at a distance of 30 km
- A Few Strategically Positioned *iGravs* will Dramatically Reduce Aliasing of Time-lapse Microgravity Surveys

Models of Volcanic Sources

➤ As the models below show - Volcanic gravity changes range in amplitude from a few μGal to several hundred μGal depending on the size and depth of the source. Changes may occur suddenly in a few seconds or slowly over days, months or years. SGs have proven their precision and stability in distinguishing hydrological models with signal amplitudes of a few tens of μGal over a similar range of periods. The *iGrav* is a proven choice to monitor and separate these complex volcanic signal sources.


Effect of a dyke and a point source as a function of the horizontal distance and:

- Two thicknesses for a dyke (case of Mt Etna, Branca et al., 2003)
- Three depths and two masses for a point source.



Precision of Spring Gravity Meters

Precision of the *iGrav*



iGrav™ Superconducting Gravity Meter

Sampling Rate: 1 Hz

Drift: Linear and $< 5 \mu\text{Gal} / \text{Year}$

Precision: $0.2 \mu\text{Gal}/\text{Hz}^{1/2} \Leftrightarrow 0.02 \mu\text{Gal} @ 100 \text{ s}$

Calibration: Stable to a Part in 10^4 for Decades!

Processes in Hydrothermal, Mid-crustal and Deep Reservoirs Beneath Volcanoes are Complex...

The Continuous, High Precision and Low Drift iGrav Superconducting Gravity Meter is Ideal to Measure and Discriminate Between These Processes.

References:

- Battaglia, M., et al., Magma intrusion beneath Long Valley Caldera confirmed by temporal changes in gravity, *Science* (285), 2119-2122, 1999.
- Battaglia, M., et al., 4D volcano gravimetry, *Geophysics*, doi:10.1190/1.2977792, 2008.
- Bonvalot, S. et al., Insights on the March 1998 eruption at Piton de la Fournaise volcano (La Réunion) from microgravity monitoring, *J. Geophys. Res.* doi:10.1029/2007JB005084, 2008.
- Branca et al., Intrusive mechanism of the 2002 NE-Rift eruption at Mt Etna (Italy) inferred through continuous microgravity data and volcanological evidences, *Geophys. Res. Lett.*, doi:10.1029/2003GL018250, 2003.
- Carbone, D., et al., A data sequence acquired at Mt Etna during the 2002-2003 eruption highlights the potential of continuous gravity observations as a tool to monitor and study active volcanoes, *J. Geodyn.*, doi:10.1016/j.jog.2006.09.012, 2006
- Carbone, D., et al., Review of microgravity observations at Mt Etna: a powerful tool to monitor and study active volcanoes, *Pure Appl. Geophys.* doi:10.1007/s00024-007-0194-7, 2007.
- Jousset et al., Temporal gravity at Merapi during the 1993-1995 crisis: an insight into the dynamical behavior of volcanoes, *J. Volcanol. Geotherm. Res.* 100, 289-320, 2000.
- Locke, C.A., et al., Magma transfer processes at persistently active volcanoes: insights from gravity observations, *J. Volcanol. Geotherm. Res.*, 127, 73-86, 2003.
- Williams-Jones, G. et al., Toward continuous 4D microgravity monitoring of volcanoes, *Geophysics*, doi:10.1190/1.2981185, 2008.
- Photographs Courtesy of WikiCommons



6264-D Ferris Square
San Diego, CA 92121
PH(858)452-7655
FAX(858)452-6965

The World's Most Stable & Precise Gravity Meter